

FIELD NOTE

Tayra (*Eira barbara*) predation of a brown-throated three-toed sloth (*Bradypus variegatus*) in Costa Rica

CAROLINA SÁENZ-BOLAÑOS^{A,1}, VÍCTOR MONTALVO^A, EDUARDO CARILLO^A
& TODD K. FULLER^B

^A Instituto Internacional en Conservación y Manejo de Vida Silvestre, Universidad Nacional, Apdo 1350–3000, Heredia, Costa Rica.
E-mail: caro.saenz0@gmail.com / csaenzb@una.cr (CSB); vmontalvog@gmail.com (VM); eduardo.carrillo.jimenez@una.cr (EC)

^B Department of Environmental Conservation, University of Massachusetts, Amherst, Massachusetts 01003, USA.
E-mail: tkfuller@eco.umass.edu

¹ Corresponding author

Abstract Being strictly arboreal, sloths become more vulnerable to predation when on the ground. Records of such predation, however, are rare. Here we present video documentation of a tayra (*Eira barbara*) preying on a juvenile brown-throated three-toed sloth (*Bradypus variegatus*) in Barbilla National Park, Costa Rica. Tayras and other ground predators, plus human activities such as habitat fragmentation, are probably major factors influencing sloth life history and permanence in the ecosystem. Predation by tayras might be more common than we think, particularly in disturbed forests.

Keywords: arboreal, Barbilla National Park, behavior, diet, prey

Depredación de perezoso de tres dedos (*Bradypus variegatus*) por tolu mucu (*Eira barbara*) en Costa Rica

Resumen Al ser estrictamente arbóreos, los perezosos se vuelven más vulnerables a la depredación cuando están en el suelo. Sin embargo, los registros de dicha depredación son raros. Aquí presentamos la documentación en video de un tolu mucu (*Eira barbara*) que caza un juvenil de perezoso de tres dedos (*Bradypus variegatus*) en el Parque Nacional Barbilla, Costa Rica. Tolu mucos y otros depredadores terrestres, además de actividades humanas como la fragmentación del hábitat, son posiblemente los principales factores que influyen en la historia de vida de los perezosos y su permanencia en los ecosistemas. La depredación por tolu mucos podría ser más común de lo que pensamos, particularmente en bosques perturbados.

Palabras clave: arbóreos, comportamiento, dieta, Parque Nacional Barbilla, presa

Of the many anti-predator options available to mammals (Caro, 2005), flight (*i.e.*, “running away”) is not one employed by sloths, often noted as being among the slowest of all living mammals (Britton, 1941b). Even though highly adapted for an arboreal lifestyle (Britton, 1941b), the slow movements and cryptic pelage of sloths do not fully protect them from canopy predators, whether avian (Beebe, 1926; Retting, 1978; Galetti & Carvalho, 2000; Touchton *et al.*, 2002; Voirin *et al.*, 2009; Aguiar-Silva *et al.*, 2017) or mammalian (Bezerra *et al.*, 2009; Sobroza *et al.*, 2016).

Even greater helplessness and vulnerability to predation of sloths on the ground (Mendel, 1985; Wainwright, 2002) is likely because “...progression is very slow [even for the sloth] and difficult” (Britton, 1941a:24). Beebe (1926:16) reported that “a mother sloth on the ground, speeded up by the calls of her infant, made 14 feet in one minute.” It is not surprising, then, that sloths spend little time on the ground. Unintended “ground-time” may result from fights resulting in falls from trees (Ingles, 1953), but sloths also come to the ground to defecate at an average interval of 8 days, with each visit lasting



FIGURE 1. Photos (from a 30-sec video) of a tayra (*Eira barbara*) preying on a juvenile brown-throated three-toed sloth (*Bradypus variegatus*) in Barbilla National Park, Costa Rica.

just a few minutes (Montgomery & Sunquist, 1975). More importantly, sloths move on the ground to get to other trees, sometimes crossing pastures (Vaughan *et al.*, 2007), roads (Britton, 1941a), and rivers (Beebe, 1926), thus exposing themselves to possible predation for longer periods of time.

The brown-throated three-toed sloth (*Bradypus variegatus*) and Hoffmann's two-toed sloth (*Choloepus hoffmanni*) have been reported as prey of several felids, including jaguars (*Panthera onca*), pumas (*Puma concolor*), and ocelots (*Leopardus pardalis*; Beebe, 1926; Garla *et al.*, 2001; Moreno *et al.*, 2006), and canids such as coyotes (*Canis latrans*) and domestic dogs (Vaughan *et al.*, 2007; Peery & Pauli, 2014). However, these observations have come mostly from studies of stomach contents, scats, or other signs; there is just a single published report of an actual predation event, specifically by a coyote of a Hoffmann's two-toed sloth on the ground (Sibaja-Morales & Cartín Núñez, 2017). Here we report the predation of a juvenile brown-throated three-toed sloth by a tayra (*Eira barbara*), known

locally as the toluuco, as documented by a video recorded by a remote camera.

During 2009–2016, we set cameras in Barbilla National Park (9°58'19"N, 83°28'51"W), located in the northern Talamanca Mountains of Costa Rica, to document the distribution and relative abundance of jaguars and other wildlife (*cf.* Sáenz-Bolaños *et al.*, 2015). At 21 locations in the park, we placed cameras (Trophy Cam models 119436, 119446, 119456, Bushnell, Overland Park, USA) on trees 0.5 m above the ground and near a human trail or animal path where we thought a jaguar might pass, and recorded either three individual photos taken 1 sec apart or a 30-sec video, with a minimum interval of 1 min between consecutive recordings.

In total, we accumulated 8,450 trap nights at the 21 camera stations. Tayras were photographed a total of 56 times at 12 (57%) of the stations; at the one station that recorded the single brown-throated three-toed sloth photographed and reported here, 14 (25%) of the tayra photos were taken. This camera trap station was located at the end of a steep

area in primary forest close to the Danta River. Forest disturbance was very low at this location; trees were big and tall with nice canopy overlap. Thus, the absence of tall trees likely was not the reason the sloth came to the ground. Moreover, the only human activity recorded in more than seven years at this station was just one video of two indigenous people with a rifle.

The video of the sloth (as best we could tell, a juvenile male) was recorded at 05:29 hr on 3 May 2013. The video begins with the tayra in the foreground walking away from the camera on a line 1 m to the left of and 2 m away from a lone sloth in the background (which the tayra is walking toward). The sloth is 0.3 m off the ground and moving down a tree perhaps 12 cm in diameter. The tayra lifts its nose, seemingly sniffing the air, walking slowly and pausing briefly. When about 1 m from the sloth, it continues sniffing but orients to the sloth (3 sec after the video begins) and then approaches it, touching it with its nose (at 8 sec; **FIG. 1**; <https://youtu.be/7WhLYIhgdeU>). The tayra sniffs both sides and underneath the sloth, the sloth moves slightly, then the tayra noses higher, swipes at the sloth briefly with its left front paw (at 17 sec), and then grabs it by its head (at 20 sec), bites around its neck and then, placing its right paw on the body, pulls it down from the tree (at 24 sec). It then appears to bite around the sloth's head and neck while the sloth weakly reaches its right arm toward the body of the tayra. The video ends at 29 sec; we believe that the sloth was killed. The next videos taken at this camera station were recorded 23 min later; they were of a tayra of the same size, apparently searching for something, sniffing all around and climbing the tree where the sloth had been. However, no sloth carcass subsequently could be found, nor could any evidence of blood, hair or something else related to the attack.

Tayras forage on the ground and in the trees of tropical and subtropical forests, and are considered opportunistic omnivores, consuming a variety of fruits, small vertebrates, insects, carrion, and honey (Presley, 2000). Tayra prey also includes medium-sized ground mammals such as rabbits (*Sylvilagus floridanus*, *S. brasiliensis*), opossums (*Didelphis marsupialis*), agoutis (*Dasyprocta punctata*), and pacas (*Agouti paca*; Galef *et al.*, 1976; Janzen, 1983; Calouro, 2000; Wainwright, 2002), and their excellent climbing skills allow them to prey on arboreal woolly opossums (*Caluromys derbianus*; Ceballos, 2016), primates such as common marmosets (*Callithrix jacchus*; Bezerra *et al.*, 2009) and red-handed howlers (*Alouatta belzebul*; Camargo & Ferrari, 2007), and sloths (Bezerra *et al.*, 2009; Sobroza *et al.*, 2016).

Sloths descend to the ground either when they are obliged to move to another tree that is not possible to reach via the canopy, or weekly to defecate. Being on the ground is a risky and energetically costly activity (Pauli *et al.*, 2014) that exposes sloths to potentially high and population-limiting predation, the latter being particularly important given the species' long lifespan and low fecundity (Peery & Pauli, 2014). Thus, abundant predators such as tayras, but including many other mesopredators, could be major factors influencing sloth life history and permanence. In addition, human activities might increase the importance of mesopredator predation on sloths. For example, habitat fragmentation could eliminate canopy connections between trees and also remove apex predators. The former might force sloths to go to the ground more frequently, while the latter might result in an increase in mesopredator abundance (Ritchie & Johnson, 2009). If so, then predation by tayras (and other mesopredators) of sloths could be more common than is currently appreciated.

ACKNOWLEDGEMENTS

We thank the National University of Costa Rica and Aventuras Naturales-Pacuare Lodge for financial and logistical support, also, the Park rangers and student assistants for help in the field, and Jim Loughry and anonymous referees for helpful comments on the manuscript.

REFERENCES

- Aguiar-Silva, F.H., O. Jaudoin, T.M. Sanaiotti, G.H.F. Seixas, S. Duleba & F.D. Martins. 2017. Camera trapping at harpy eagle nests: interspecies interactions under predation risk. *Journal of Raptor Research* 51: 72–78. <https://doi.org/10.3356/JRR-15-58.1>
- Beebe, W. 1926. The three-toed sloth *Bradypus cuculliger* Wagler. *Zoologica* 7: 1–67.
- Bezerra, B.M., A. Barnett, A. Souto & G. Jones. 2009. Predation by the tayra on the common marmoset and the pale-throated three-toed sloth. *Journal of Ethology* 27: 91–96. <https://doi.org/10.1007/s10164-008-0090-3>
- Britton, S.W. 1941a. Form and function in the sloth. *The Quarterly Review of Biology* 16: 13–34. <https://doi.org/10.1086/394620>
- Britton, S.W. 1941b. Form and function in the sloth (concluded). *The Quarterly Review of Biology* 16: 190–207. <https://doi.org/10.1086/394628>
- Calouro, A.M. 2000. Attempted predation on Brazilian rabbit (*Sylvilagus brasiliensis* — Lagomorpha: Leporidae) by tayra (*Eira barbara* — Carnivora: Procyonidae). *Revista de Biología Tropical* 48: 267–268.
- Camargo, C.C. & S.F. Ferrari. 2007. Interactions between tayras (*Eira barbara*) and red-handed howlers (*Alouatta belzebul*) in eastern Amazonia. *Primates* 48: 147–150. <https://doi.org/10.1007/s10329-006-0009-y>

- Caro, T. 2005. Antipredator defenses in birds and mammals. University of Chicago Press, Chicago, Illinois, USA. 592 pp.
- Ceballos, G. 2016. Observation of attempted predation of a woolly opossum (*Caluromys derbianus*) by a tayra (*Eira barbara*) during daylight in Belize. *Revista Mexicana de Mastozoología Nueva Época* 6: 46–49.
- Galef, B.G. Jr., R.A. Mittermeier & R.C. Bailey. 1976. Predation by the tayra (*Eira barbara*). *Journal of Mammalogy* 57: 760–761. <https://doi.org/10.2307/1379450>
- Galetti, M. & O. Carvalho, Jr. 2000. Sloths in the diet of a harpy eagle nestling in Eastern Amazon. *The Wilson Bulletin* 112: 535–536. [https://doi.org/10.1676/0043-5643\(2000\)112\[0535:SITDOA\]2.0.CO;2](https://doi.org/10.1676/0043-5643(2000)112[0535:SITDOA]2.0.CO;2)
- Garla, R.C., E.Z. Setz & N. Gobbi. 2001. Jaguar (*Panthera onca*) food habits in Atlantic rain forest of southeastern Brazil. *Biotropica* 33: 691–696.
- Ingles, L.G. 1953. Observations on Barro Colorado Island mammals. *Journal of Mammalogy* 34: 266–268. <https://doi.org/10.1093/jmammal/34.2.266a>
- Janzen, D.H. 1983. Costa Rican natural history. University of Chicago Press, Chicago, Illinois, USA. 816 pp.
- Mendel, F.C. 1985. Use of hands and feet of three-toed sloths (*Bradypus variegatus*) during climbing and terrestrial locomotion. *Journal of Mammalogy* 66: 359–366. <https://doi.org/10.2307/1381249>
- Montgomery, G.G. & M.E. Sunquist. 1975. Impact of sloths on Neotropical forest energy flow and nutrient cycling. Pp. 69–98 in: *Tropical ecological systems* (F.B. Golley & E. Medina, eds.). Springer Verlag, New York. https://doi.org/10.1007/978-3-642-88533-4_7
- Moreno, R.S., R.W. Kays & R. Samudio, Jr. 2006. Competitive release in diets of ocelot (*Leopardus pardalis*) and puma (*Puma concolor*) after jaguar (*Panthera onca*) decline. *Journal of Mammalogy* 87: 808–816. <https://doi.org/10.1644/05-MAMM-A-360R2.1>
- Pauli, J.N., J.E. Mendoza, S.A. Steffan, C.C. Carey, P.J. Weimer & M.Z. Peery. 2014. A syndrome of mutualism reinforces the lifestyle of a sloth. *Proceedings of the Royal Society B* 281: 20133006. <https://doi.org/10.1098/rspb.2013.3006>
- Peery, M.Z. & J.N. Pauli. 2014. Shade-grown cacao supports a self-sustaining population of two-toed but not three-toed sloths. *Journal of Applied Ecology* 51: 162–170. <https://doi.org/10.1111/1365-2664.12182>
- Presley, S.J. 2000. *Eira barbara*. *Mammalian Species* 636: 1–6. <https://doi.org/10.2307/0.636.1>
- Retting, N.L. 1978. Breeding behavior of the harpy eagle (*Harpia harpyja*). *The Auk* 95: 629–643.
- Ritchie, E.G. & C.N. Johnson. 2009. Predator interactions, mesopredator release and biodiversity conservation. *Ecology Letters* 12: 982–998. <https://doi.org/10.1111/j.1461-0248.2009.01347.x>
- Sáenz-Bolaños, C., V. Montalvo, T.K. Fuller & E. Carrillo. 2015. Records of black jaguars at Parque Nacional Barbilla. *Cat News* 62: 38–39.
- Sibaja-Morales, K.D. & M. Cartín Núñez. 2017. Unusual predation of Hoffmann's two-toed sloth (*Choloepus hoffmanni*) by a coyote (*Canis latrans*) in Costa Rica. *Edentata* 18: 89–91. <https://doi.org/10.2305/IUCN.CH.2017.Edentata-18-1.13.en>
- Sobroza, T.V., A.L. Gonçalves & L. Silva dos Santos. 2016. Predation attempt and abnormal coat coloration of the tayra (*Eira barbara*) in the Brazilian Central Amazon. *Studies on Neotropical Fauna and Environment* 51: 231–234. <https://doi.org/10.1080/01650521.2016.1227137>
- Touchton, J., Y.C. Hsu & A. Palleroni. 2002. Foraging ecology of reintroduced captive-bred subadult harpy eagles (*Harpia harpyja*) on Barro Colorado Island, Panama. *Ornitología Neotropical* 13: 365–379.
- Vaughan, C., O. Ramírez, G. Herrera & R. Guries. 2007. Spatial ecology and conservation of two sloth species in a cacao landscape in Limon, Costa Rica. *Biodiversity and Conservation* 16: 2293–2310. <https://doi.org/10.1007/s10531-007-9191-5>
- Voirin, B., R. Kays, M. Lowman & M. Wikelski. 2009. Evidence of three-toed sloth (*Bradypus variegatus*) predation by spectacled owl (*Pulsatrix perspicillata*). *Edentata* 8–10: 15–20. <https://doi.org/10.1896/020.010.0113>
- Wainwright, M. 2002. The natural history of Costa Rican mammals. Zona Tropical Publications, Miami, Florida, USA. 383 pp.

Received: 2 June 2018; Accepted: 26 July 2018